

SCIENTIFIC LETTER

Length of the free tendon is not associated with return to play time in biceps femoris muscle injuries



La longitud del tendón libre no está asociada al tiempo de vuelta a la competición en las lesiones musculares del bíceps femoral

Injuries of the hamstring muscle complex are very common injuries in sports, representing 12% of all injuries and 37% of all muscle injuries in professional football¹ with biceps femoris as the most frequently (84%) injured hamstring muscle.²

The long head of the biceps femoris originates on the posterior aspect of the ischial tuberosity. It shares a footprint and (portion of its) tendon with the semitendinosus.^{3–8} The tendon can be subdivided into a 'free' and 'intramuscular' tendon (sometimes referred to as 'central tendon'^{9–11}). The free tendon is defined as the part of the tendon that has no muscle fibers inserting onto it, the remaining part of the tendon is referred to as the intramuscular tendon.

In multi-sports club Futbol Club Barcelona (FC Barcelona), anecdotal clinical observations led to the clinical suspicion that a short proximal free tendon in injuries of the long head of the biceps femoris (BFLH) was associated with a quicker return to play (RTP). Measurement reliability and potential prognostic value (i.e. time to return to play) of free proximal tendon length have not yet been evaluated. Our hypothesis is that a shorter proximal BFLH free tendon is associated with a shorter time to return to play (RTP).

Therefore, the primary aim of the present study was:

- To evaluate the reliability of the proximal BFLH free tendon measurement on magnetic resonance imaging (MRI).
- To evaluate the association between proximal BFLH free tendon length and time to RTP in days.

Methods

Subjects

To address our research question, a study population comprised of two different study cohorts was used (50 Catalan cohort and 58 Qatari cohort). Patients with acute hamstring injuries were either recruited at the multi-sports club FC Barcelona, or were part of a double-blind randomized controlled trial (RCT) investigating the effect of platelet-rich plasma (PRP) in acute hamstring injury on time to RTP.¹³ This RCT did not show a benefit of PRP over placebo. Eligibility criteria for both study cohorts are shown in [Box 1](#).

Ethical approval was obtained from the Ethical Committee of ASPETAR (Orthopaedic and Sports Medicine Hospital) and the Consell Català de l'Esport (Generalitat de Catalunya). Informed consent was obtained from all included subjects.

MRI protocol

At FC Barcelona, images were acquired with a 3.0 T magnet system (Siemens, Erlangen, Germany). Subjects were positioned inside the scanner with both legs parallel to the MRI table, and the thighs were covered with multichannel coils. Coronal and axial T1-weighted images (TR/TE 900/15 ms; FOV 330 mm × 300 mm; slice thickness 3.5 mm; matrix 384 × 512) were obtained. Coronal and axial T2 fat sat (TR/TE 4000/35 ms; FOV 330 mm × 300 mm; slice thickness 3.5 mm; matrix 320 × 384) as well as sagittal T2 (TR/TE 4000/32 ms; FOV 330 mm × 300 mm; slice thickness 2.0 mm; matrix 320 × 384) was obtained.

In the ASPETAR cohort, images were acquired with a 1.5 T magnet system (Magnetom Espree, Siemens) and a body matrix coil. Coronal and axial proton density (PD) weighted images (TR/TE 3000/32 ms; FOV 240 mm; slice thickness 5 mm; matrix, 333 × 512) were obtained. Subsequently, coronal and axial proton density weighted images with fat

Box 1		
Study cohort	Catalan cohort	Qatari cohort (Hamilton et al. ¹²)
Inclusion criteria	<ul style="list-style-type: none"> • Acute onset of posterior thigh pain • Localized pain on palpation • Localized pain of passive stretching • Increased pain on isometric contraction • MRI confirmed a grade I or II BFLH lesion 	<ul style="list-style-type: none"> • Age 18–50 years • Available for follow-up • Acute onset of posterior thigh pain • Presenting an MRI within 5 days from injury • MRI confirmed a grade I or II BFLH lesion • Male gender • Able to perform five sessions of physiotherapy a week at our clinic
Exclusion criteria	<ul style="list-style-type: none"> • Contraindication to MRI • Chronic hamstring injury • Concurrent other injury inhibiting rehabilitation 	<ul style="list-style-type: none"> • Contraindication to MRI • Reinjury or chronic hamstring injury • Concurrent other injury inhibiting rehabilitation • Unwilling to comply with follow-up • Needle phobia • Overlying skin infection • Diabetes, immunocompromised state • Medication with increasing bleeding risk • Medical contraindication to injection




Figure 1 Proximal BFLH free tendon length is measured (in cm) from the most inferior margin of the ischial tuberosity to the point where the first muscle fibers start to insert onto the tendon. IT: ischial tuberosity; ST: semitendinosus; BFLH: long head of the biceps femoris; *: free tendon. (a) Short free tendon, (b) medium free tendon and (c) long free tendon.

saturation (PD-FS) were obtained (TR/TE 3000/32 ms; FOV 240 mm; slice thickness 3.5 mm; matrix 326 × 512 for coronal and TR/TE 3490/27 ms; FOV 320 mm; slice thickness 3.5 mm; matrix 333 × 512 for axial).

MRI assessments

Each MRI was assessed by one of the two musculoskeletal radiologists, who were blinded to clinical outcome. Each radiologist scored the MRIs in a random order between July 2009 and July 2016. The free proximal BFLH tendon measurement is shown in [Figs. 1 and 2](#).




Figure 2 T1 coronal MRI of a short free tendon (a) and long free tendon (b) are shown. Arrowheads show the most inferior margin of the ischial tuberosity and arrows show the point where the first muscle fibers started to insert onto the tendon.

Rehabilitation protocol

The rehabilitation protocols used in both cohorts are described in detail in previous studies.^{13,14}

Outcome measures

At FC Barcelona, the RTP decision is made based on clinical assessment, sport-specific field tests and imaging.^{15,16} The field test includes a comparison of data such as workload parameters (including GPS) with athlete-specific pre-injury data. No specific pre-defined criteria were used. Time to RTP was defined as the number of days from initial injury




Figure 3 Flow diagram of the inclusion process.

until the athlete was cleared by one of the team physicians to resume full, unrestricted training.

In the ASPETAR cohort, the guidelines for making the final RTP decision included successful and asymptomatic completion of the progressive criteria-based rehabilitation program including three sport-specific phases, clinical evaluation by an experienced sports medicine physician, and interpretation of the results of an isokinetic assessment. The physician's final decision was guided, but not determined by these medical factors, and also included consideration of sport risk modifiers and decision modifiers.^{17,18}

Statistical analysis

Statistical analysis was carried out using IBM SPSS Statistics (version 21, IBM Corp.). Continuous variables were tested for normality and presented as mean (\pm SD) unless otherwise stated. To evaluate intra-observer reliability, one of two specialized musculoskeletal radiologists recorded free tendon length bilaterally in 30 subjects. The same radiologist repeated the measurements one week later. To evaluate inter-observer reliability, both radiologists performed the measurements independently. Intraclass correlation coefficient (ICC) analysis was carried out to determine intra- and inter-observer reliability. We used Cronbach's α and the model for the computation of ICC was a two-way mixed (people effects are random, and the item effects are fixed). Reliability was considered excellent if the ICC is >0.75 , fair to good if $0.4 < \text{ICC} < 0.75$ and poor if ICC is <0.4 . The Pearson correlation coefficient was used to analyze the relationship between tendon length (in cm) and time to RTP (in days). Statistical significance was set at $p < 0.05$.

Results

From 2008 to 2014, 108 athletes with a BFLH injury were included for analysis (Fig. 3).

Demographics and injury characteristics at baseline are shown in Table 1.

Reliability

ICCs for intra-observer and inter-observer reliability of the free tendon length measurement (cm) are presented in Table 2.

Free tendon length and RTP time

Average length of the free proximal BFLH tendon was 4.8 ± 2.4 cm on the right side, and 5.0 ± 2.2 cm on the left

Table 1 Baseline characteristics of the included subjects.

	N = 108
<i>Age (years)</i>	25 (± 5.8)
<i>Sports category</i>	
Athletics	4 (4%)
Baseball	1 (1%)
Basketball	7 (7%)
Decathlon	1 (1%)
Field hockey	7 (7%)
Football	63 (58%)
Futsal	7 (7%)
Handball	2 (2%)
Physical coach	1 (1%)
Track and field	13 (12%)
Volleyball	1 (1%)
Weight lifting/body building	1 (1%)
<i>Side injured</i>	
Right	68 (63%)
Left	40 (37%)
Bilateral	0 (0.0%)
<i>Injury location</i>	
Proximal	78 (72%)
Distal	30 (28%)
<i>Anatomic location</i>	
Free tendon	0 (0%)
Musculotendinous junction	104 (96%)
Myofascial	4 (4%)
<i>Free tendon length (BFLH)</i>	
Right free tendon length (cm)	4.8 (± 2.4)
Left free tendon length (cm)	5.0 (± 2.2)


side. Average time to RTP was 29 ± 15 days. There was no statistically significant correlation between free proximal BFLH tendon length in cm and time to RTP in days (Pearson correlation coefficient = -0.037 ; $p = 0.712$). Fig. 4 shows a scatter plot for these parameters.

Discussion

In the present study, we sought to determine whether or not a relationship exists between proximal BFLH free tendon length and RTP time, a hypothesis that was formed based on clinical observations. The presented measurement of proximal BFLH free tendon length has excellent intra- and

Table 2 Intraclass correlation coefficients (ICC).

Reliability	Tendon	ICC	95% CI	p-Value	Strength of agreement
Intra-observer	Right	0.983	0.963–0.992	<.001	Excellent
	Left	0.921	0.832–0.963	<.001	Excellent
Inter-observer	Right	0.984	0.966–0.993	<.001	Excellent
	Left	0.977	0.952–0.989	<.001	Excellent

**Figure 4** Scatter plot for RTP (days) and free proximal BFLH tendon length (cm).

inter-observer reliability. However, there was no correlation between tendon length (cm) and RTP time (days).

Long head biceps femoris proximal free tendon and RTP

Athletes in the current study are predominantly football players and track and field athletes with a mean time to RTP of approximately 4 weeks. For comparison: a large study on RTP in (elite) football reported that 84% of injuries involved the biceps femoris, with a mean time to RTP of 20 ± 15 days.¹⁹ Askling et al.²⁰ studied 18 sprinters who all had the primary injury in the biceps femoris with a median time to return to pre-injury activity level of around 16 weeks (range 6–50 weeks).

According to our measurements, average proximal length of BFLH free tendon was 4.8 ± 2.4 cm on the right side and 5.0 ± 2.2 cm on the left side. A recent anatomical study of the hamstring muscle complex included muscle and tendon lengths and reported an average proximal BFLH free tendon length of 5.0 ± 3.4 cm.⁵ Although this study involved

measurements upon dissection, the average of this tendon length corresponds well with our measurements on MRI.

The clinical suspicion that proximal BFLH free tendon length might impact recovery was partly due to observations in clinical practice, and partly due to recent attention for the effect of musculotendinous architecture on injury risk.^{21–23} Hypothetically, if certain architectural characteristics result in conditions that may predispose to muscle injury, they could also potentially result in less favorable conditions for muscle recovery. We sought to investigate whether this was the case for proximal free tendon length. Nonetheless, our data do not indicate that free proximal BFLH tendon length influences time to RTP following acute hamstring injury in athletes.

In any case, is interesting to note that tendinous injuries (for example proximal BFLH free tendon tears) could need a surgical intervention for a whole healing because in case of perform a conservative treatment the RTP is much longer and the risk of reinjury are high.²⁴ However, a recent work²⁵ has pointed that RTP in intramuscular hamstring tendon

injuries is only significantly longer compared with injuries without intramuscular tendon disruption.

Limitations

The subjects in this study were either participants in a randomized controlled trial (Qatari cohort) or athletes of FC Barcelona (Catalan cohort) with some differences in eligibility criteria, MRI protocols and RTP criteria. Although this may reflect clinical practice, it can be considered as a limitation. In addition, a univariate analysis was used. Therefore, we have not controlled for potential confounders.

Future research

Aside from impact on recovery, in accordance with recent studies on architectural characteristics it would be interesting to investigate whether free proximal BFLH tendon length is a risk factor for injury and re-injury. This requires prospective study designs.

Conclusion

Free tendon length can be reliably measured with excellent intra-observer and inter-observer agreement. In long head biceps femoris injury, length of the proximal free tendon is not associated with time to return to play.

Conflict of interests

The authors declare that that they have no competing interests.

References

- Ekstrand J, Hägglund M, Kristenson K, Magnusson H, Waldén M. Fewer ligament injuries but no preventive effect on muscle injuries and severe injuries: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med.* 2013;47:732–7.
- Ekstrand J, Healy JC, Walden M, Lee JC, English B, Hägglund M. Hamstring muscle injuries in professional football: the correlation of MRI findings with return to play. *Br J Sports Med.* 2012;46:112–7.
- Battermann N, Appell H-J, Dargel J, Koebke J. An anatomical study of the proximal hamstring muscle complex to elucidate muscle strains in this region. *Int J Sports Med.* 2011;32:211–5.
- Feucht MJ, Plath JE, Seppel G, Hinterwimmer S, Imhoff AB, Brucker PU. Gross anatomical and dimensional characteristics of the proximal hamstring origin. *Knee Surg Sports Traumatol Arthrosc.* 2015;23:2576–82.
- van der Made AD, Wieldraaijer T, Kerkhoffs GM, Kleipool RP, Engebretsen L, van Dijk CN, et al. The hamstring muscle complex. *Knee Surg Sports Traumatol Arthrosc.* 2015;23:2115–22.
- Philippson MJ, Ferro FP, Campbell KJ, Michalski MP, Goldsmith MT, Devitt BM, et al. A qualitative and quantitative analysis of the attachment sites of the proximal hamstrings. *Knee Surg Sports Traumatol Arthrosc.* 2015;23:2554–61.
- Sato K, Nimura A, Yamaguchi K, Akita K. Anatomical study of the proximal origin of hamstring muscles. *J Orthop Sci.* 2012;17:614–8.
- Woodley SJ, Mercer SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs.* 2005;179:125–41.
- Comin J, Malliaras P, Baquie P, Barbour T, Connell D. Return to competitive play after hamstring injuries involving disruption of the central tendon. *Am J Sports Med.* 2013;41:111–5.
- Pollock N, James SL, Lee JC, Chakraverty R. British athletics muscle injury classification: a new grading system. *Br J Sports Med.* 2014;48:1347–51.
- van der Made A, Almusa E, Whiteley R, Hamilton B, Eirale C, van Hellemond F, et al. Intramuscular tendon involvement on MRI has limited value for predicting time to return to play following acute hamstring injury. *Br J Sports Med.* 2018.
- Hamilton B, Tol JL, Almusa E, Boukarroum S, Eirale C, Farooq A, et al. Platelet-rich plasma does not enhance return to play in hamstring injuries: a randomised controlled trial. *Br J Sports Med.* 2015;49:943–50.
- Tol JL, Hamilton B, Eirale C, Muxart P, Jacobsen P, Whiteley R. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficits. *Br J Sports Med.* 2014;48:9–1364.
- Valle X, Tol L, Hamilton J, Rodas B, Malliaras G, Malliaropoulos P, et al. *Asian J Sports Med.* 2015;6:e12541.
- Serveis. *Mèdics Futbol Club Barcelona Clinical Practice Guide for muscular injuries. Epidemiology, diagnosis, treatment and prevention. Version 4.5 (9 February 2009). Apunts Med Esport.* 2009;164:179–203.
- Pruna R, Andersen TE, Clarsen B, McCall A. *Muscle injury guide: prevention of and return to play from muscle injuries.* Barcelona: Ed Barça Innovation Hub; 2018.
- Creighton DW, Shrier I, Shultz R, Meeuwisse WH, Matheson GO. Return-to-play in sport: a decision-based model. *Clin J Sport Med.* 2010;20:379–85.
- Tol JL, Hamilton B, Eirale C, Muxart P, Jacobsen P, Whiteley R. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficits. *Br J Sports Med.* 2014;48:1364–9.
- Ekstrand J, Lee JC, Healy JC. MRI findings and return to play in football: a prospective analysis of 255 hamstring injuries in the UEFA Elite Club Injury Study. *Br J Sports Med.* 2016;50:738–43.
- Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med.* 2006;35:197–206.
- Evangelidis PE, Massey GJ, Pain MTG, Folland JP. Biceps femoris aponeurosis size: a potential risk factor for strain injury? *Med Sci Sports Exerc.* 2015;47:1383–9.
- Fiorentino NM, Blemker SS. Musculotendon variability influences tissue strains experienced by the biceps femoris long head muscle during high-speed running. *J Biomech.* 2014;47:3325–33.
- Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C, Opar DA. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med.* 2016;50:1524–35.
- Yanguas J, Pruna R, Puigdemívol J, Mechó S. Clinical and imaging aspects of assessment and management of proximal long head biceps femoris injury (free-tendon and miotendinosus junction injuries). A report of two cases. *Apunts Med Esport.* 2017;52:79–82.
- van der Made AD, Almusa E, Reurink G, Whiteley R, Weir A, Hamilton B, et al. Intramuscular tendon injury is not associated with an increased hamstring reinjury rate within 12 months after return to play. *Br J Sports Med.* 2018;52:1261–6.

Ricard Pruna^a, Javier Yanguas^{a,*},
Anne D. van der Made^{b,c,d}, Lluís Capdevila^e,
Ramon Balius^f, Xavier Alomar^g, Javier Arnaiz^b,
Johannes L. Tol^{b,c,d}, Gil Rodas^a

^a *Medical Department Futbol Club Barcelona, FIFA Medical Center of Excellence, Barcelona, Spain*

^b *Aspetar, Orthopedic and Sports Medicine Hospital, Doha, Qatar*

^c *Academic Center for Evidence Based Sports Medicine, Academic Medical Center, University of Amsterdam, Amsterdam Movement Sciences, Amsterdam, Netherlands*

^d *Department of Orthopaedic Surgery, Academic Medical Center, University of Amsterdam, Amsterdam Movement Sciences, Amsterdam, Netherlands*

^e *Laboratory of Sport Psychology, Universitat Autònoma de Barcelona, Barcelona, Spain*

^f *Consell Català de l'Esport, Generalitat de Catalunya, Barcelona, Spain*

^g *Radiology Department, Centres Mèdics Creu Blanca, Barcelona, Spain*

* Corresponding author.

E-mail address: xavier.yanguas@fcbarcelona.cat

(J. Yanguas).

Available online 22 January 2019