

ORIGINAL ARTICLE

# Could regular practice of volleyball modulate salivary secretory immunity in children? Cross-sectional and longitudinal studies



Carolina da Silva Peres<sup>a</sup>, Roberta Pratti Gava<sup>a</sup>, Natália Miwa Yoshida<sup>b</sup>,  
Julio Cesar Molina Correa<sup>c</sup>, Lorena Beatriz Scudeller<sup>b</sup>, Gabriela Fleury Seixas<sup>d</sup>,  
Carla Cristiane Silva<sup>e,\*</sup>, Cassia Cilene Dezan Garbelini<sup>f</sup>, Solange de Paula Ramos<sup>g</sup>

<sup>a</sup> Health Sciences Center, State University of Londrina, Rodovia Celso Garcia Cid, PR 445 km380, Londrina-PR, Brazil

<sup>b</sup> State University of Londrina, Rodovia Celso Garcia Cid, PR 445 km380, Londrina-PR, Brazil

<sup>c</sup> Center of Physical Education and Sports, State University of Londrina, Rodovia Celso Garcia Cid, PR 445 km380, Londrina-PR, Brazil

<sup>d</sup> Faculty of Odontology, Universidade Norte do Paraná, Rua Marselha, 60 Londrina-PR, Brazil

<sup>e</sup> Department of Physical Education, State University of the North of Paraná, Alameda Padre Magno, 841 Jacarezinho, Londrina-PR, Brazil

<sup>f</sup> Department of Oral Medicine and Pediatrics Dentistry, Health Sciences Center, State University of Londrina, Rodovia Celso Garcia Cid, PR 445 km380, Londrina-PR, Brazil

<sup>g</sup> Department of Histology, Center of Biological Sciences, State University of Londrina, Rodovia Celso Garcia Cid, PR 445 km380, Londrina-PR, Brazil

Received 30 July 2019; accepted 4 February 2020

Available online 24 February 2020

## KEYWORDS

Secretory immunoglobulin A;  
Cardiorespiratory fitness;  
Muscle strength;  
Youth sports

## Abstract

**Introduction:** The aim of these studies was to evaluate the effects of regular volleyball training on salivary SIgA and alpha-amylase in female children and adolescents.

**Material and method:** In the cross-sectional study, 115 female children ( $12.6 \pm 2.2$  years) participating in a Volleyball training program were classified as beginners, intermediate, and competitive level. The children were evaluated regarding caries index, body mass index (BMI), cardiorespiratory fitness, and countermovement jump. In the longitudinal study: 54 girls (intermediate and competitive groups) trained for 8 weeks and were re-evaluated at weeks 4 and 8.

**Results:** The SIgA secretion rate and alpha-amylase activity did not present correlations with training category, age, caries index, or training status. A weak positive correlation was detected between IgA secretion rate and BMI ( $r=0.20$ ,  $p<0.05$ ). After 8 weeks of training,  $VO_2\max$  ( $p<0.005$ ) and jump height ( $p<0.005$ ) improved in the competitive girls. No differences were observed in salivary secretion rate, SIgA concentration and secretion rate, or alpha-amylase activity at weeks 4 and 8.

\* Corresponding author.

E-mail address: [ccsilva@uenp.edu.br](mailto:ccsilva@uenp.edu.br) (C.C. Silva).

**Conclusion:** We concluded that despite improving physical fitness, 8-weeks of recreational volleyball training are not able to improve salivary SIgA secretion or alter alpha-amylase activity in female children and adolescents.

## Introduction

The practice of regular physical exercises at moderate intensity or recreational sports training is associated with positive modulation of secretory immune responses and decreased stress in adult populations.<sup>1-5</sup> Salivary secretory Immunoglobulin A (SIgA) is the most abundant antibody in saliva and plays important roles in the defense of the oral cavity against dental caries, periodontal diseases, and oral and upper respiratory tract infections.<sup>5-7</sup> Clinical data have suggested that regular exercise practice and increased physical fitness were correlated to increased SIgA levels and improved oral health.<sup>1,3,8,9</sup> In this way, improving physical fitness may have positive effects on oral immunity.

Salivary SIgA secretion is driven by the autonomic nervous system and its secretion can be modulated by physical exercise.<sup>5,10,11</sup> A single bout of moderate exercise (75% VO<sub>2</sub>max) is able to improve oral mucosal immunity over 24 h, increasing salivary secretion of SIgA.<sup>12</sup> Other authors demonstrated that aerobic exercise training programs could increase resting SIgA secretion in adults.<sup>3,4</sup> A study in female undergraduate students demonstrated that students engaged in regular team sports training (in season) presented lower levels of stress markers and increased levels of SIgA compared to the "off season" students.<sup>1</sup> While acute and regular moderate exercise can improve salivary secretion, low levels of physical activity are associated with impairments in the immune response in adults.<sup>2,5</sup> However, little is known about the effects of regular exercise practice on salivary secretion in children and adolescents.

Salivary alpha amylase is a digestive and antimicrobial enzyme that has been used to monitor the stress balance and activation of the sympathetic nervous system.<sup>13,14</sup> Some studies have suggested that amylase levels are modulated by stress, body mass index (BMI), and physical fitness in children and adults.<sup>1,15,16</sup> In adult men, muscle power and cardiovascular fitness are associated with better regulation of the autonomic nervous system and lower secretion of salivary amylase.<sup>15</sup> Female undergraduate students engaged in regular team sport training demonstrated reduced levels of stress and salivary alpha-amylase secretion.<sup>1</sup> A low cardiorespiratory fitness level was also associated with increased sympathetic activity and salivary alpha-amylase concentration in school children.<sup>16</sup> Higher levels of alpha-amylase are associated with increased BMI and emotional stress in pre-school children.<sup>11,16</sup> However, other authors did not find differences in salivary alpha-amylase concentration between normal and overweight children aged 5–12 years.<sup>18</sup> Currently, it is not known whether physical training could modulate resting alpha-amylase secretion, since it may improve autonomic nervous system modulation and body mass index control.<sup>14,19</sup>

Although the benefits of regular physical activity have been reported in adult and elderly populations,<sup>3,4,20,21</sup> the effects of regular practice of exercise on salivary secre-

tory immunity in children and adolescents are not yet clear. A study in African children reported that overweight and obese children presented decreased cardiorespiratory fitness, lower salivary SIgA concentration and secretion rate, and increased levels of salivary alpha-amylase.<sup>16</sup> Since engagement in physical activities improves immunity defenses in adult populations,<sup>3,5-21</sup> it may also have some beneficial effects on children and adolescents. Therefore, the objectives of the present studies were to determine whether engagement of female children and adolescents in volleyball training could improve secretory immunity in saliva and the effects of changing physical performance on SIgA and alpha-amylase secretion.

## Material and methods

### Study design

This study was divided into two steps:

- Cross-sectional study: a transversal study on the association between saliva flow rate, SIgA concentration, SIgA secretion rate, and alpha-amylase concentration with BMI and physical fitness;
- Cohort study: a longitudinal study of the effects of 8 weeks of volleyball training on salivary secretory immunity.

The study was approved by the Research Ethics Committee Involving Human Beings of the State University of Londrina (protocol no. 821.804) and followed the principles of the Helsinki Declaration (2013).<sup>22</sup> All parents or legal guardians, as well as the children and adolescents, gave their written informed consent.

In the cross-sectional study, 148 girls were included, ranging in age from 7 to 17 years. The girls trained in the Project of School Sports of the Department of Sports of the Municipality of Cambé – Paraná, Brazil. The samples were collected during a previous study (da Silva et al., 2019)<sup>23</sup> of our research group that evaluated their salivary inflammatory profile. The girls were evaluated in the first month of training (August, 2015) after the July (2015) vacation period. During the vacation period (30 days), girls and parents reported no participation in systematic training and sports activities. In the initial evaluation, we included all the girls who presented a free and informed consent signed by the parents and themselves. We excluded girls who presented chronic inflammatory diseases, oral mucosal lesions, continued use of medication, diabetics and other chronic inflammatory diseases, bronchitis and asthma, recent use of corticosteroids (less than 6 months), and those who donated an insufficient sample volume for amylase analysis and could not perform the physical tests.

In the cohort study, the girls were re-evaluated after four and eight weeks of training. Only girls who donated a sufficient volume of saliva and were regularly attending training sessions were included in the final evaluation, with up to two absences in training sessions allowed during the investigation period. In addition, a control group of 40 girls ( $12.4 \pm 2.17$  years, normal weight) from local schools, not participating in sports training, were evaluated for salivary SIgA and amylase at baseline and after 8 weeks.

## Participants

The initial evaluation (Pre) was used in the cross-sectional study analysis. The girls were submitted to physical evaluation, saliva sampling, the countermovement jump test (CMJ), and 20 m shuttle run test (Léger test). Regarding their previous participation in the School Sports Project, the girls were classified into three categories:

- **Beginners** (without previous experience): the girls had no previous experience in volleyball practice and were not engaged in other sports activities. These girls had less than 3 weeks of training.
- **Intermediate group**: minimum time of three months and maximum of 12 months of engagement in volleyball training. The girls had participated in the sports program since the previous semester, but did not participate regularly in competitive events.
- **Competitive group**: minimum of 13 months of training. The girls in the competitive group had participated in volleyball training for more than two school semesters and were engaged in local and state school game competitions.

## Saliva sampling

The saliva samples were collected as previously described by da Silva et al.<sup>23</sup> The saliva samples were collected at rest before physical tests and training commenced, 1 h after their last meal, between 09:00 and 10:00 a.m. and 2:00 p.m. and 3:00 p.m., according to the training schedule. The girls were asked to brush their teeth 1 h before saliva sampling to eliminate excess toothpaste residues from saliva. The girls were instructed to rinse their mouths with drinking water for 1 min prior to saliva collection, and then salivate spontaneously into sterile graduated collection tubes for 2 min. The salivary flow rate was assessed by measuring the volume of secreted saliva per minute (ml/min). Samples were frozen at  $-20^{\circ}\text{C}$  prior to use. Menstrual phase was not controlled because it has been demonstrated not to influence secretion of SIgA and other antimicrobial proteins at rest.<sup>24</sup> Unstimulated whole saliva was used since stimulated salivation could increase saliva flow rate and alpha amylase secretion or cause bias in the SIgA secretion rate calculus.<sup>25</sup>

## Enzyme-linked immunosorbent assay for determination of salivary IgA

The saliva samples were centrifuged at  $4000 \times g$  for 5 min for sedimentation of cell debris. Supernatants from the saliva

samples were diluted at 1:1000 in phosphate buffered saline (PBS, pH 7.2). Next, the samples were submitted to an ELISA assay for determination of the salivary SIgA concentration, using commercial kits as recommended by the manufacturer (A88-102P, Bethyl Laboratories, Montgomery, USA). The SIgA concentration was expressed in  $\mu\text{g/ml}$  saliva and the secretion rate was determined by IgA secretion per minute ( $\mu\text{g/min}$ ).<sup>26</sup>

## Detection of saliva alpha-amylase

The concentration of salivary alpha-amylase was determined with a commercial enzyme kit (Phadebas Pharmacia Diagnostica, Uppsala, Sweden), as recommended by the manufacturer.

## Physical evaluation

Height in centimeters was measured using a stadiometer and weight was evaluated using a digital scale (Omron HBF 514C, Omron Health Care do Brasil, São Paulo, Brazil). The body mass index was calculated using the formula  $\text{BMI} = \text{mass (kg)}/\text{height}^2 \text{ (m)}$ . The girls were classified according to BMI into: very underweight, underweight, normal, overweight, and obese according to the criteria adopted by the World Health Organization for girls between 0 and 19 years of age.<sup>27</sup>

The oral examination was performed by one trained Periodontist and two Pediatric dentists, ( $\text{Kappa} = 0.99$ ). An oral examination was performed using artificial light, spatulas, dental mirrors, and probes. The index of decayed, missing, and filled teeth for permanent teeth (DMFT) was determined according to the criteria described by the World Health Organization.<sup>28</sup> During the oral examination, the presence of soft tissue lesions and spontaneous gingival bleeding were also recorded.

## 20 m shuttle run test (Léger test)

The maximum oxygen consumption ( $\text{VO}_2\text{max}$ ) was estimated using the maximum progressive test developed by Léger and Lambert in 1982 and modified by Léger et al. in 1984.<sup>29</sup> The formula described by Léger et al. in 1988 for children aged 8–19 years<sup>30</sup> was used to calculate estimated cardiorespiratory fitness.

## Countermovement jump test

The girls were familiarized with the countermovement jump test prior to data collection. The vertical jumping test with countermovement was performed on a contact mat (Smart jump, Fusion Sports, Summer Park, Australia),<sup>31</sup> with three consecutive jumps with a 1 min interval. The highest jump height in three attempts was recorded. For the test, the children were instructed to stand erect and, when a light signal came on, crouch quickly with their knees bent at approximately  $90^{\circ}$  and immediately jump with their legs extended. The arms were extended during the flight phase of the jump, simulating the blocking movement at the net. The phase of

menstrual cycle was not controlled since it does not influence muscular strength in young females.<sup>32</sup>

### Volleyball training program

The volleyball training practices were performed in two to three sessions per week, lasting 120–180 min daily, depending on their category and previous experience in sports. The training was performed for eight consecutive weeks, and consisted of 10 min of warm-up, specific technical-tactical exercises for the category, aerobic and anaerobic exercises with and without balls, simulated games, and 5 min of stretching at the end of each training session. Children who were absent three or more times during the eight weeks of training were excluded from the study.

The girls evaluated in the longitudinal test completed the eight weeks of training and were submitted to the Léger test and vertical jump test in the first (Pre) and eighth weeks (Post). Saliva collections were performed at the beginning (Pre) and after four (4 weeks) and eight weeks (Post) of training.

### Statistical analysis

The distribution of normality was evaluated by the Shapiro–Wilks test. Continuous variables with normal distribution are expressed as mean and standard deviation. Variables without normal distribution are expressed as medians and quartiles of 25–75%. Differences between groups were assessed using the one-way ANOVA test, with the Tukey post-test (parametric data) or Kruskal–Wallis test (non-parametric data). Categorical data are expressed in absolute frequencies and percentages and were evaluated using the Chi-square test with the Yates correction, or Fisher's exact test. The correlations between the salivary measurements and other variables were determined by the Spearman's rank-order correlation test. Two-way ANOVA was performed in the cohort study to determine the influence of time and previous experience in volleyball practice on physical performance and salivary secretion. The sphericity of data was analyzed by Mauchly's test with the Greenhouse–Geisser correction. Differences between study variables were considered significant at  $p < 0.05$ . Statistical analysis was performed with Graph Pad Prism 8.0 (GraphPad software, San Diego, USA).

We considered a 50% difference in the mean SIgA concentration reported by Starzak et al.,<sup>16</sup> for normal weight ( $243.0 \pm 199.2 \mu\text{g/ml}$ ) children, for sample size calculation. Twenty-two children per group was necessary to achieve a maximum  $\alpha$  error of 5%, and 80% statistical power (BioEstat 5.0, Instituto de Desenvolvimento Sustentável Mamirauá, Tefe, Brazil).

## Results

### Cross-sectional study

In total, 148 girls were enrolled in the volleyball program in August 2015, of which 115 (77.7%) were evaluated; 33 (22.3%) were excluded due to lack of presentation of

the consent form, absence on the day of physical testing, or insufficient volume of saliva sample to perform alpha-amylase analysis.

The age ranged from 7 years to 17 years. Girls in the competitive group were older than the other groups (Table 1). The frequency of girls in different BMI ranges was similar between groups ( $p > 0.05$ ; Chi squared test).

The mean  $\text{VO}_2\text{max}$  was not significantly different between groups. However, the competitive girls presented a higher jump height compared to the beginners (Table 1).

No significant differences in saliva flow rate, SIgA concentration, SIgA secretion rate, or alpha amylase activity were detected between groups (Table 1).

The saliva flow rate, SIgA concentration and secretion rate, and alpha-amylase activity did not present correlation with age, caries index (DMFT), or physical performance (Table 2). A week correlation was observed between IgA secretion rate and BMI score (Table 2).

### Longitudinal study

Only 54 (46.9%) girls completed the assessments and had enough volume of saliva samples for analysis after 8 weeks of training. Three girls (05.5%) were from the beginner group, 22 (40.7%) from the intermediate group, and 29 (53.7%) from the competitive group. Considering that 24 (87.5%) girls in the beginner group did not complete the training sessions or perform the physical tests, this group was excluded from the longitudinal study.

The girls in the intermediate group did not present improvement in  $\text{VO}_2\text{max}$  ( $p = 0.58$ , paired  $t$  test; Fig. 1a) or jump height ( $p = 0.35$ , paired  $t$  test; Fig. 1b). On the other hand, girls from the competitive group presented improvement in both  $\text{VO}_2\text{max}$  ( $p = 0.003$ , paired  $t$  test, Fig. 1a) and jump height ( $p = 0.002$ , paired  $t$  test, Fig. 1b). An effect was observed in aerobic performance only for time ( $p = 0.009$ ;  $F = 7.58$ , two-way ANOVA). Jump performance presented group ( $p = 0.002$ ,  $F = 10.13$ ) and time ( $p = 0.008$ ,  $F = 7.74$ , two-way ANOVA) interactions.

The levels of saliva flow rate, IgA concentration and secretion rate, and alpha amylase activity did not change significantly from Pre values to weeks 4 and Post (Fig. 1). No group or time effects and interactions were observed in either group in any salivary measurement.

The salivary variables of the beginners and competitive group were not different from a control group at pre and post ( $p > 0.05$ , Kruskal–Wallis test). For the control group, the variations from pre to post for saliva flow rate were ( $1.19 \pm 1.23$  to  $0.73 \pm 0.41 \text{ ml/min}$ ,  $p < 0.05$ , Wilcoxon test), IgA concentration ( $133.6 \pm 18.3$  to  $169.7 \pm 46.3 \mu\text{g/ml}$ , Wilcoxon test), IgA secretion rate ( $159.1 \pm 21.7$  to  $124.3 \pm 18.9 \mu\text{g/min}$ ,  $p < 0.05$ , Paired  $t$  test), and amylase ( $18.1 \pm 3.0$  to  $18.5 \pm 2.8 \text{ mmol/ml}$ ,  $p < 0.05$ , paired  $t$  test).

## Discussion

Contrary to our hypothesis, physical fitness was not associated with regulation of salivary secretion of SIgA and alpha-amylase in female children and adolescents performing volleyball training. Improvement in physical fitness was

**Table 1** Characteristics of participants and salivary IgA and alpha amylase in girls with different training groups.

	Beginners N=24	Intermediate N=52	Competitive N=39	Total N=115
Age (years)	11.8 ± 2.3	12.3 ± 2.0	13.4 ± 2.2 <sup>**</sup> ,#	12.5 ± 2.5
DMFT	1 [0.5–1.5]	1 [0.5–2.5]	1 [0.5–3.0]	1 [0.5–2.5]
<b>BMI</b>				
Underweight	1 (4.1%)	5 (9.6%)	1 (2.5%)	7 (6.1%)
Normal	17 (70.8%)	35 (67.3%)	31 (79.5%)	83 (72.1%)
Overweight	3 (12.5%)	5 (9.6%)	5 (12.8%)	13 (11.3%)
Obese	3 (12.5%)	7 (13.4%)	2 (5.1%)	12 (10.5%)
Number of weekly training sessions	–	2 [2–2.5]	2 [2–3.0]	2 [2–2.5]
Total time of weekly training (min)	–	240 [210–300]	300 [240–540] <sup>#</sup>	240 [210–300]
VO <sub>2</sub> max (ml/kg/min)	39.4 ± 5.9	40.0 ± 4.3	39.2 ± 5.1	39.7 ± 4.9
Jump height (cm)	24.8 ± 3.9	28.1 ± 6.0	30.8 ± 5.2 <sup>**</sup>	28.4 ± 5.7
Saliva flow rate (ml/min)	0.70 ± 0.40	0.70 ± 0.30	0.70 ± 0.40	0.70 ± 0.40
SlgA concentration (µg/ml)	181.4 [141.0–262.4]	179.9 [138.7–308.6]	193.4 [133.2–350.1]	188.8 [135.8–291.5]
SlgA secretion rate (µg/min)	142.6 [76.7–231.4]	130.4 [65.8–203.4]	126.5 [57.4–206.0]	134.4 [65.8–203.4]
Alpha Amylase activity (µmol/s*l)	17.5 [15.8–20.2]	17.5 [16.5–18.9]	17.6 [16.8–18.8]	17.5 [16.0–18.9]

DMFT: decayed, missing and filled teeth (caries index); BMI: body mass index; VO<sub>2</sub>max: maximum oxygen consumption, SlgA: salivary immunoglobulin A.

<sup>\*\*</sup> p < 0.01, competitive vs. beginners.

<sup>#</sup> p < 0.05, competitive vs. intermediate.

**Table 2** Correlation between saliva flow rate, SlgA concentration and secretion rate, and Alpha amylase activity with physical characteristics.

	Saliva flow rate		IgA concentration		IgA secretion rate		Alpha Amylase activity	
	r	p value	r	p value	r	p value	r	p value
Age	0.09	0.31	0.07	0.42	0.09	0.30	0.11	0.20
DMFT	–0.03	0.74	–0.14	0.13	–0.14	0.13	0.07	0.42
BMI	0.10	0.25	0.17	0.08	0.20	0.03*	0.05	0.56
CMJ	0.04	0.71	0.13	0.24	0.06	0.60	0.19	0.06
VO <sub>2</sub> max	0.09	0.31	0.00	0.99	0.06	0.60	0.10	0.25
SlgA (µg/ml)	0.17	0.06	–	–	–	–	0.07	0.42
SlgA (µg/min)	–	–	–	–	–	–	0.01	0.92
Alpha Amylase	0.09	0.31	0.07	0.42	0.01	0.92	–	–


DMFT: decayed, missing and filled teeth (caries index); BMI: body mass index; VO<sub>2</sub>max: maximum oxygen consumption, SlgA: salivary immunoglobulin A.

more evident in the competitive group but was not associated with improved secretion of SlgA or downmodulation of alpha-amylase activity. This was surprising; since other studies reported salivary secretion and immunity were modulated by physical training in adults.<sup>1,4,5,20</sup> Moreover, a previous study in young female subjects suggested that regular sports training could improve secretory immunity and reduce alpha-amylase secretion.<sup>1</sup> We excluded two girls with active caries from the study since these could increase the salivary levels of SlgA and alpha-amylase.<sup>33</sup> The results demonstrated that girls presented similar caries activity between study groups and the caries index was not associated with variability in SlgA and alpha-amylase secretion.

Previous studies investigating salivary levels of SlgA in school children and adolescents presented divergent results.<sup>16,34,35</sup> One study investigated cellular and SlgA immune responses in children (aged 7–13 years) corre-

lated with skinfold thickness, BMI, and VO<sub>2</sub>peak.<sup>35</sup> The authors concluded that increased adiposity was correlated with impaired cellular responses but increased SlgA secretion, whereas aerobic fitness presented little effect on immune parameters.<sup>35</sup> One study reported that obesity and low cardiorespiratory fitness were associated with reduced levels of salivary SlgA and increased alpha-amylase activity in children aged 10 ± 1.6 years.<sup>16</sup> However, another study demonstrated that SlgA concentration was increased in obese/overweight children (aged 6–12 years) and presented a weak correlation with BMI.<sup>34</sup> A limitation of studies, including the current study, is missing data on maturation status and a wide variation in age (children to adolescents). A previous study investigated the relationship between age (8–17 years old), hormonal status, and grip strength with salivary IgA.<sup>36</sup> As observed in the present study, no correlations were detected with age and strength, but a moderate correlation with dehydroepiandrosterone was





**Figure 1** Median and 25–75% interquartile range of VO<sub>2</sub>max (a), counter movement jump height (b), saliva flow rate (c), salivary SlgA concentration (d), salivary SlgA secretion rate, (e) and alpha-amylase activity (f). \*\*\* $p < 0.05$ , difference in competitive group from Pre to Post, Wilcoxon test.

observed.<sup>36</sup> This suggests that other maturation parameters such as hormonal status, menarche, and puberty should be investigated in future studies addressing secretory immunity in female children and adolescents. Another concern regards the menstrual cycle phase, since some reports in adult women suggested that the association between menstrual phase and sexual activity may have some impact on salivary SlgA secretion,<sup>37,38</sup> while others found no correlation of menstrual cycle with salivary secretory immunity.<sup>24,39</sup> We expected a lower secretion rate of SlgA in overweight and obese children, since increased BMI is associated with reduced parasympathetic activity in children and adolescents.<sup>40</sup> Contrary to our hypothesis, only a weak positive correlation was observed between SlgA secretion rate and BMI. The lower limb strength and VO<sub>2</sub>max were also not correlated with resting SlgA and alpha amylase activity. The low number of obese and overweight children observed in our study may be a limitation of the study. Another concern is that children were detrained at the initial evaluation (cross-sectional study), so training effects on mucosal immunity could be blunted. However, 8-weeks of training improved physical performance, without a significant effect on saliva secretion. Moreover, children who were not training presented similar levels of salivary SlgA and alpha-amylase and did not present significant changes in salivary secretion after 8 weeks.

In the cross-sectional study, the competitive girls presented increased jump height. This difference may be associated with differences in the age of the groups since increased maturational status was positively associated with jumping performance in youth athletes.<sup>41</sup> Another hypothesis was that the girls from the intermediate and competitive groups were evaluated after a period of detraining (school vacations), although lower limb strength adaptations are maintained for longer periods than aerobic adaptations.<sup>42</sup> During the training period, jump and aerobic performance significantly increased in the competitive girls, suggesting a previous state of adaptation to exercise. In young soccer athletes, a brief period of detraining impaired both strength and endurance performances, although a fast recovery in performance was observed after training began.<sup>43</sup> Furthermore, the competitive group presented higher weekly training volume than the intermediate group, which may account for the increased physical performance in comparison to the intermediate group after 8 weeks. Indeed, increased training volume may be necessary in adolescents (>13 years) to achieve the same level of strength performance improvements as younger children.<sup>44</sup> However, we could not monitor individual training load during the training period to draw any correlations among training volume, physical adaptation, and immune parameters. Unfortunately, controlling training load by heart rate or the

lactate threshold was not feasible in this study due to the large number of participants. Using rate of perceived exertion to calculate internal training load was not appropriate as it is influenced by trainability, body mass index, and age.<sup>45,46</sup> Thus, we made the assumption that competitive children had a higher training load due to a higher training volume, considering that the technical–tactical activities developed by both groups were similar. As expected, the girls engaged in competition events and who had trained for a longer period presented better performance improvement. However, despite the improvements in performance, no effect of training stimulus was observed on salivary secretory immunity.

A study in 10 year-old Norwegian children revealed that practice of team sports, including volleyball, was associated with increased cardiorespiratory fitness.<sup>47</sup> Other authors demonstrated that 7 weeks of Volleyball training in female adolescent athletes improved aerobic fitness and vertical jump height.<sup>48</sup> In male child and adolescent volleyball athletes, increased anaerobic power was associated with increased vertical jump height, compared to nonathletic subjects.<sup>49</sup> Although improvements were observed in physical performance, the physiological stimulus seemed not to be sufficient to increase resting levels of SIgA and alpha-amylase in female children and adolescents. Indeed, aerobic performance was the main component altered by the volleyball training and a previous study demonstrated that aerobic fitness was not correlated with SIgA.<sup>35</sup>

Human studies have demonstrated that physical training and increased cardiorespiratory fitness could improve humoral immunity and are associated with improved parasympathetic balance.<sup>1,3,16,20</sup> Moreover, increased parasympathetic activity can induce SIgA secretion, stimulating poli-Ig receptor translocation of secretory immunoglobulins into saliva.<sup>11,50</sup> We hypothesize that during the initial evaluation, cardiorespiratory fitness was not significantly different between groups and this may account for the lack of significant differences in autonomic regulation of SIgA secretion. Indeed, increased  $\text{VO}_2\text{peak}$  and  $\text{VO}_2\text{max}$  were associated with better overall regulation of the autonomic nervous system in children and adolescents.<sup>51,52</sup> Moreover, engagement in sports activity above 180 min/week was associated with improved vagal autonomic regulation in children aged 10–13 years,<sup>53</sup> so improved cardiovascular fitness may have some positive impact on salivary secretion in children and adolescents due to improved autonomic regulation. Considering the lack of differences in  $\text{VO}_2\text{max}$  between groups after returning from school vacations (a long detraining period), no differences in SIgA secretion were expected. However, although during the 8 weeks of volleyball training,  $\text{VO}_2\text{max}$  significantly increased in the competitive group, this was not accompanied by changes in salivary secretion. A recent systematic review concluded that physical training may have little impact on autonomic regulation in healthy children<sup>54</sup> and may not account for changes in salivary secretion. Therefore, the improved aerobic capacity may not reflect a positive impact on parasympathetic regulation of salivary secretion.

Previous studies have suggested that increased BMI may account for increased levels of alpha-amylase and decreased salivary SIgA in healthy children.<sup>16,17</sup> Increased BMI is associ-

ated with low cardiorespiratory fitness and poor autonomic regulation, although these effects can be overcome by physical training in adult individuals.<sup>55</sup> In the present study, BMI was not correlated with alpha amylase activity, suggesting other factors may regulate saliva secretion in this study population. Salivary secretion of alpha-amylase is modulated by parasympathetic–sympathetic outflow on salivary glands and is considered a marker of sympathetic modulation,<sup>56,57</sup> which could potentially be modulated by training status and fitness.<sup>12,15</sup> Contrary to our hypothesis, no correlation was found between fitness and improvement in performance with secretion of salivary alpha-amylase at rest. In older men, salivary alpha amylase response is similar in low and highly physically active subjects, suggesting no differences in this salivary biomarker at rest and without stressful stimulus.<sup>58</sup> In the same way, it is possible that no difference in alpha amylase activity was detected in the present study because the girls were not exposed to a stressful challenge during saliva sampling and might not have upregulated sympathetic outflow under training adaptation.

We concluded that despite improving physical fitness, 8 weeks of recreational volleyball training is not able to improve salivary SIgA secretion or alter alpha-amylase activity in female children and adolescents. In the studied population, physical fitness and the caries index were not related to SIgA and alpha-amylase levels.

## Conflict of interest

The authors declare that they don't have any conflict of interest.

## Acknowledgements

The authors would like to thank the Coordination for the Improvement of Higher Education Personnel (CAPES) for granting a scholarship to C.P.S. (grant number 06172286986/2015) and the Fundação Araucária – PR for granting a scholarship to R.G.P. (174/2015). We thank the National Council for Research and Development (CNPq) for granting funding for the development of the study (grant no. 482524/2013-8). We thank the Secretary of Sports of the Municipality of Cambé, Paraná, Brazil and the coordinating teachers of School Volleyball, Reginaldo Mazzola and Eduardo Fiel.

## References

1. Lamb AL, Hess DE, Edenborn S, Ubinger E, Carrillo AE, Appasamy PM. Elevated salivary IgA, decreased anxiety, and an altered oral microbiota are associated with active participation on an undergraduate athletic team. *Physiol Behav.* 2017;169:169–77.
2. Trochimiak T, Hubner-Wozniak E. Effect of exercise on the level of immunoglobulin a in saliva. *Biol Sport.* 2012;29:255–61.
3. Sloan CA, Engels HJ, Fahlman MM, Yarandi HE, Davis JE. Effects of exercise on S-IGA and URS in postmenopausal women. *Int J Sports Med.* 2013;34:81–6.
4. Martins RA, Cunha MR, Neves AP, Martins M, Teixeira-Verissimo M, Teixeira AM. Effects of aerobic conditioning on salivary IgA and plasma IgA IgG and IgM in older men and women. *Int J Sports Med.* 2009;30:906–12.

5. Bishop NC, Gleeson M. Acute and chronic effects of exercise on markers of mucosal immunity. *Front Biosci (Landmark Ed)*. 2009;14:4444–56.
6. Colombo NH, Pereira JA, da Silva ME, Ribas LF, Parisotto TM, Mattos-Graner Rde O, et al. Relationship between the IgA antibody response against *Streptococcus mutans* GbpB and severity of dental caries in childhood. *Arch Oral Biol*. 2016;67:22–7.
7. Giuca MR, Pasini M, Tecco S, Giuca G, Marzo G. Levels of salivary immunoglobulins and periodontal evaluation in smoking patients. *BMC Immunol*. 2014;15:5.
8. Ferreira RO, Correa MG, Magno MB, Almeida A, Fagundes NCF, Rosing CK, et al. Physical activity reduces the prevalence of periodontal disease: systematic review and meta-analysis. *Front Physiol*. 2019;10:234.
9. Anderson AP, Park YM, Shrestha D, Zhang J, Liu J, Merchant AT. Cross-sectional association of physical activity and periodontal antibodies. *J Periodontol*. 2018;89:1400–6.
10. Carpenter GH, Proctor GB, Garrett JR. Preganglionic parasympathectomy decreases salivary SIgA secretion rates from the rat submandibular gland. *J Neuroimmunol*. 2005;160:4–11.
11. Proctor GB, Carpenter GH. Salivary secretion: mechanism and neural regulation. *Monogr Oral Sci*. 2014;24:14–29.
12. Rosa L, Teixeira A, Lira F, Tufik S, Mello M, Santos R. Moderate acute exercise (70% VO<sub>2</sub> peak) induces TGF- $\beta$ ,  $\alpha$ -amylase and IgA in saliva during recovery. *Oral Dis*. 2014;20:186–90.
13. Adam EK, Till Hoyt L, Granger DA. Diurnal  $\alpha$  amylase patterns in adolescents: associations with puberty and momentary mood states. *Biol Psychol*. 2011;88:170–3.
14. Strahler J, Skoluda N, Kappert MB, Nater UM. Simultaneous measurement of salivary cortisol and  $\alpha$ -amylase: application and recommendations. *Neurosci Biobehav Rev*. 2017;83:657–77.
15. Wyss T, Boesch M, Roos L, Tschopp C, Frei KM, Annen H, et al. Aerobic fitness level affects cardiovascular and salivary  $\alpha$  amylase responses to acute psychosocial stress. *Sports Med Open*. 2016;2:33.
16. Starzak DE, Konkol KF, McKune AJ. Effects of cardiorespiratory fitness and obesity on salivary secretory IgA and  $\alpha$ -amylase in South African children. *Children (Basel)*. 2016:3.
17. Miller AL, Sturza J, Rosenblum K, Vazquez DM, Kaciroti N, Lumeng JC. Salivary  $\alpha$  amylase diurnal pattern and stress response are associated with body mass index in low-income preschool-aged children. *Psychoneuroendocrinology*. 2015;53:40–8.
18. de Campos MM, Kobayashi FY, Barbosa Tde S, Costa Sda S, Lucas Bde L, Castelo PM. Characteristics of salivary secretion in normal-weight, overweight and obese children: a preliminary study: salivary composition and excessive fat tissue. *Odontology*. 2014;102:318–24.
19. Oliveira RS, Barker AR, Wilkinson KM, Abbott RA, Williams CA. Is cardiac autonomic function associated with cardiorespiratory fitness and physical activity in children and adolescents? A systematic review of cross-sectional studies. *Int J Cardiol*. 2017;236:113–22.
20. de Araujo AL, Silva LC, Fernandes JR, Matias Mde S, Boas LS, Machado CM, et al. Elderly men with moderate and intense training lifestyle present sustained higher antibody responses to influenza vaccine. *Age (Dordr)*. 2015;37:105.
21. Lee PH. Physical activity, sedentary behaviors, and Epstein-Barr virus antibodies in young adults. *Physiol Behav*. 2016;164:390–4.
22. World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310:2191–4.
23. da Silva CP, Pratti RG, Molina JCC, Silva CCd, Garbelini CCD, Ramos SP. Salivary levels of Interleukin-6 and Tumor Necrosis Factor- $\alpha$  in girls aged 7 to 17 years practicing volleyball. *Apunts Med IEsport*.
24. Gillum T, Kuennen M, Miller T, Riley L. The effects of exercise, sex, and menstrual phase on salivary antimicrobial proteins. *Exerc Immunol Rev*. 2014;20:23–38.
25. Allgrove JE, Oliveira M, Gleeson M. Stimulating whole saliva affects the response of antimicrobial proteins to exercise. *Scand J Med Sci Sports*. 2014;24:649–55.
26. Milanez VF, Ramos SP, Okuno NM, Bousso DA, Nakamura FY. Evidence of a non-linear dose-response relationship between training load and stress markers in elite female futsal players. *J Sports Sci Med*. 2014;13:22–9.
27. de Onis M, Lobstein T. Defining obesity risk status in the general childhood population: which cut-offs should we use? *Int J Pediatr Obes*. 2010;5:458–60.
28. WHO. Oral health surveys: basic methods. 5th ed. WHO; 2013. p. 125.
29. Leger LA, Lambert J. A maximal multistage 20-m shuttle run test to predict VO<sub>2</sub> max. *Eur J Appl Physiol Occup Physiol*. 1982;49:1–12.
30. Leger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci*. 1988;6:93–101.
31. Reeve TC, Tyler CJ. The validity of the SmartJump contact mat. *J Strength Cond Res*. 2013;27:1597–601.
32. Arazi H, Nasiri S, Eghbali E. Is there a difference toward strength, muscular endurance, anaerobic power and hormonal changes between the three phase of the menstrual cycle of active girls? *Apunts Med IEsport*. 2019;54:65–72.
33. Sharma A, Subramaniam P, Moiden S. Analysis of salivary IgA,  $\alpha$ -amylase, lactoferrin, and lysozyme before and after comprehensive dental treatment in children: a prospective study. *Contemp Clin Dent*. 2017;8:526–30.
34. Perez MM, Pessoa JS, Ciamponi AL, Diniz MB, Santos M, Alves HHO, et al. Correlation of salivary immunoglobulin A with Body Mass Index and fat percentage in overweight/obese children. *J Appl Oral Sci*. 2018;27:e20180088.
35. Nieman DC, Henson DA, Fagoaga OR, Nehlsen-Cannarella SL, Sonnenfeld G, Utter AC. Influence of skinfold sum and peak VO<sub>2</sub> on immune function in children. *Int J Obes Relat Metab Disord*. 2002;26:822–9.
36. Hodges-Simeon CR, Prall SP, Blackwell AD, Gurven M, Gaulin SJC. Adrenal maturation, nutritional status, and mucosal immunity in Bolivian youth. *Am J Hum Biol*. 2017:29.
37. Lorenz TK, Demas GE, Heiman JR. Interaction of menstrual cycle phase and sexual activity predicts mucosal and systemic humoral immunity in healthy women. *Physiol Behav*. 2015;152:92–8.
38. Lorenz TK, Heiman JR, Demas GE. Testosterone and immune-reproductive tradeoffs in healthy women. *Horm Behav*. 2017;88:122–30.
39. Burrows M, Bird SR, Bishop N. The menstrual cycle and its effect on the immune status of female endurance runners. *J Sports Sci*. 2002;20:339–44.
40. Eyre EL, Duncan MJ, Birch SL, Fisher JP. The influence of age and weight status on cardiac autonomic control in healthy children: a review. *Auton Neurosci*. 2014;186:8–21.
41. Asadi A, Ramirez-Campillo R, Arazi H, Saez de Villarreal E. The effects of maturation on jumping ability and sprint adaptations to plyometric training in youth soccer players. *J Sports Sci*. 2018;36:2405–11.
42. Lo MS, Lin LL, Yao WJ, Ma MC. Training and detraining effects of the resistance vs. endurance program on body composition, body size, and physical performance in young men. *J Strength Cond Res*. 2011;25:2246–54.
43. Chatzinikolaou A, Michaloglou K, Avloniti A, Leontsini D, Deli CK, Vlachopoulos D, et al. The trainability of adolescent soccer



- players to brief periodized complex training. *Int J Sports Physiol Perform.* 2018;13:645–55.
44. Rodriguez-Rosell D, Franco-Marquez F, Mora-Custodio R, Gonzalez-Badillo JJ. Effect of high-speed strength training on physical performance in young soccer players of different ages. *J Strength Cond Res.* 2017;31:2498–508.
  45. Abe D, Yoshida T, Ueoka H, Sugiyama K, Fukuoka Y. Relationship between perceived exertion and blood lactate concentrations during incremental running test in young females. *BMC Sports Sci Med Rehabil.* 2015;7:5.
  46. Tompkins CL, Flanagan T, Lavoie J 2nd, Brock DW. Heart rate and perceived exertion in healthy weight and obese children during a self-selected physical activity program. *J Phys Act Health.* 2015;12:976–81.
  47. Resaland GK, Aadland E, Andersen JR, Bartholomew JB, Andersen SA, Moe VF. Physical activity preferences of 10-year-old children and identified activities with positive and negative associations to cardiorespiratory fitness. *Acta Paediatr.* 2018;108:354–60.
  48. Eliakim A, Portal S, Zadik Z, Meckel Y, Nemet D. Training reduces catabolic and inflammatory response to a single practice in female volleyball players. *J Strength Cond Res.* 2013;27:3110–5.
  49. Kasabalis A, Douda H, Tokmakidis SP. Relationship between anaerobic power and jumping of selected male volleyball players of different ages. *Percept Mot Skills.* 2005;100:607–14.
  50. Kurimoto Y, Saruta J, To M, Yamamoto Y, Kimura K, Tsukinoki K. Voluntary exercise increases IgA concentration and polymeric Ig receptor expression in the rat submandibular gland. *Biosci Biotechnol Biochem.* 2016;80:2490–6.
  51. Tanha T, Wollmer P, Fedorowski A, Thorsson O, Karlsson MK, Dencker M. Correlation between physical activity, aerobic fitness and body fat against autonomic function profile in children. *Clin Auton Res.* 2016;26:197–203.
  52. da Silva DF, Bianchini JA, Antonini VD, Hermoso DA, Lopera CA, Pagan BG, et al. Parasympathetic cardiac activity is associated with cardiorespiratory fitness in overweight and obese adolescents. *Pediatr Cardiol.* 2014;35:684–90.
  53. Radtke T, Khattab K, Brugger N, Eser P, Saner H, Wilhelm M. High-volume sports club participation and autonomic nervous system activity in children. *Eur J Clin Invest.* 2013;43:821–8.
  54. da Silva CC, Pereira LM, Cardoso JR, Moore JP, Nakamura FY. The effect of physical training on heart rate variability in healthy children: a systematic review with meta-analysis. *Pediatr Exerc Sci.* 2014;26:147–58.
  55. Tian Y, Huang C, He Z, Hong P, Zhao J. Autonomic function responses to training: correlation with body composition changes. *Physiol Behav.* 2015;151:308–13.
  56. Nater UM, Rohleder N. Salivary alpha-amylase as a non-invasive biomarker for the sympathetic nervous system: current state of research. *Psychoneuroendocrinology.* 2009;34:486–96.
  57. Bosch JA, Veerman EC, de Geus EJ, Proctor GB. alpha-Amylase as a reliable and convenient measure of sympathetic activity: don't start salivating just yet! *Psychoneuroendocrinology.* 2011;36:449–53.
  58. Strahler J, Fuchs R, Nater UM, Klaperski S. Impact of physical fitness on salivary stress markers in sedentary to low-active young to middle-aged men. *Psychoneuroendocrinology.* 2016;68:14–9.