

ORIGINAL ARTICLE

Observational study with the objective of determining possible correlations between GRF and muscle activation at reception after a jump in an ACL injury



Alesander Badiola-Zabala^{a,*}, Nuria Massó-Ortigosa^a, Josep Cabedo-Sanromà^b, Fernando Rey-Abella^a, Raimón Milà^a, Guillermo Ruben Oviedo^{a,b}

^a *Facultat de Ciències de la Salut Blanquerna, Universitat Ramon Llull, Barcelona, Spain*

^b *Facultat de Psicologia, Ciències de l'Educació i de l'Esport, BLanquerna Universitat Ramon Llull, Barcelona, Spain*

Received 18 January 2020; accepted 31 March 2020

Available online 2 June 2020

KEYWORDS

Anterior cruciate ligament (ACL);
Ground reaction forces (GRF);
Landing;
Neuromuscular activation

Abstract

Introduction: The ACL injury is considered one of the most serious injuries and usually occurs in actions that include movements with changes of direction, jump and landing. It is a common injury between the young active population and the risk in women of suffering from non-contact injury is superior to that of men. Athletes who suffer from non-contact injuries of the ACL usually have common biomechanical profiles, with landings with large values in ground reaction force (GRF) and therefore, low cushioning on landing.

Objective: To determine possible correlations between GRF and muscular activation at landing after a jump.

Material and methods: The type of study carried out is an observational study in which, using surface electromyography (EMG), a force platform and an electrogoniometer, the aim is to assess muscle activation and its relationship with GRF (specifically the vertical component Fz).

Results: Correlations have been observed between the reaction force of the soil (Fz) in the moments where the reaction force of the soil is greater and the instant where the knee reaches maximum flexion after landing, with the activation of certain muscle groups and differences depending on the gender of the subject.

Discussion: The neuromuscular recruitment strategies in the phases of maximum GRF load and knee flexion are different depending on the sex of the individual, so it should be considered when scheduling prevention and recovery work.

Conclusion: The evaluation of GRF and muscle activation patterns, allows to assess the dynamics of landing after a jump and to be able to detect different patterns according to sex, with the consequent importance that it can have in the injury mechanism.

* Corresponding author.

E-mail address: alesanderbz@blanquerna.url.edu (A. Badiola-Zabala).

Introduction

Knee injuries and specifically ACL injuries are prevalent in a wide variety of activities (competitive or not) and especially in those that include acceleration-deceleration movements, changes of direction, jump and landing. They are common among young active population¹ and the risk in women of suffering the non-contact injury is higher than that in men,²⁻¹⁰ twice,¹ five times^{11,12} and up to six or seven times more^{13,14} or greater. Athletes who suffer from non-contact injury of the ACL, usually have common biomechanical profiles, with landings with large values in GRF and therefore, little cushioning at landing.^{15,16}

In the field of sports, ACL injury is one of the most common and most serious knee injuries, along with meniscal injury.¹⁷⁻²² Although the actual incidence of ACL injuries is not fully known, it can be estimated through ACL reconstructions that are performed a year before an injury, these are placed on a fork between 200,000 and 350,000 a year in the USA alone²³⁻²⁹ and in 38 out of every 100,000 inhabitants in Denmark,³⁰ one third affecting women.²⁸

The ACL injury is multifactorial (neuromuscular, biomechanical, anatomical, genetic, hormonal, etc.). Although it has been studied, little is known about why the structural weakness of the ligament. The importance of this aspect is that the external forces acting on the tissues also depend on the intrinsic properties of each. And the hormonal aspect can modify these structural properties making it vulnerable to external loads. Therefore, risk factors may also differ depending on gender.³¹

In the analysis of a jump, the landing phase is of vital importance and a clear example of them is that 31% of the injuries occur on the landing after a jump,^{32,33} so it is a moment of activity that deserves to be thoroughly studied.

Descriptive studies of non-contact ACL injury indicate that the injury occurs shortly after initial landing contact or after deceleration, with minimal or no contact in 70% of cases.^{32,34,35}

In a normal landing pattern, the muscles contract to absorb the forces of GRF (ground reaction forces) but there may be an abnormal absorption of GRF^{32,36} and an example of this is that the ischiocrural muscle activity during landing it can vary between 20% and 60% of the value of the maximum voluntary contraction.³⁷

During landing, hip, knee and ankle actions help absorbing reaction forces from the ground. When these segments are not effective in the synergistic work of absorbing GRF, the leg acts as a column of two segments, which sometimes makes it unable to absorb the force of GRF.³⁸ In this same landing activity after a jump, the traction and torsion forces are increased in the ACL with the increase of the ground reaction forces (GRF) and the decrease of the hip angles^{38,39} and one more straight position, it is associated with an increase in GRF.^{38,40,41} A landing with a more flexed trunk and an increase in hip and knee flexion can reduce GRF.^{38,40,42}

In relation to muscle strength, a reduction, as well as a reduction in muscle preactivity prior to contact with the ground during sports activities, are factors related with an increased risk of ACL injury in female athletes. Maximum voluntary strength and pre-activation do not have

a direct relationship and therefore a person with little muscle strength does not necessarily have to have a bad pre-activation. In order for this pre-activation to be effective, these muscles have to co-activate quickly and adequately since the latency of the feedback sensor reflex is greater than 75–100 ms.⁴³⁻⁴⁵

Pre-activation is considered important because the ACL lesion occurs 40ms after contact of the foot with the ground at reception after a jump (the GRF peak occurs 40ms after contact with the ground)⁴⁵⁻⁵¹ being the axial compression force of the knee during landing, 6 times the body weight^{52,53} and the reason for this may be that a power imbalance between external forces and muscle contractions may partly explain why the injuries of ACL does not usually occur in the window of less than 40ms. Since the foot contacts the ground.^{43,51,56} Some authors extend this period of time, indicating the range between 30ms and 100ms of the initial contact of the foot on the ground.^{28,46,51,52}

The posterior kinetic chain muscles are especially important: gluteus maximus and medius, hamstrings, and gastrocnemius and soleus. The muscles of the posterior chain must be recruited correctly in order to absorb the GRF; otherwise it will be the ligaments and the joint who should do it. The gluteus maximus, the second strongest muscle in the human body, is the only tri-axial plane controller of the femoral position. When an athlete primarily contracts his quadriceps and reduces the contraction of the gluteus and hamstrings, the result is the collapse of the knee in valgus. And this can be problematic because this allows the GRF to act in a valgus position⁵⁴ and epidemiological studies indicate that a large knee abduction moment, large knee abduction angles, and the GRF are important predictors of ACL injury risk with 78% sensitivity and 73% specificity.^{52,55}

Neuromechanically, reflex learning regulates joint stiffness and it has been suggested that these mechanisms are centrally pre-programmed (CNS)⁵⁶⁻⁵⁹ and it is not clear whether these pre-programmed activation patterns can be voluntarily changed. Voluntary changes at this level are difficult to achieve.^{56,60} However, on the external or internal focus of attention, changes in the landing mechanisms have been proven, suggesting that a conscious adaptation of control of pre-programmed landing patterns could be possible.^{56,60,61} From the neuromechanical point of view, joint stabilization is achieved during voluntary movements with compensatory and anticipatory adjustments to minimize unpredictable and predictable disturbances.⁶¹⁻⁶⁴ For this reason, we can observe abnormal responses to predictable and unpredictable disturbances in ACL patients, years after the injury; This could mean functional instability during activities of daily living^{61,65-67} and therefore, an increased risk of re-injury.^{61,68}

For this reason, it is considered important to evaluate the relationship between GRF and the activation of different muscle groups during landing after a jump and see if sex can be a variable that influences these activation patterns. This information can be useful for the evaluation of risk factors, as well as for the programming of activities with preventive purposes in specific populations.

Material and methods

Material

- Biopac model MPC 150 was used to capture the biomedical signals.
- For the recording of the electromyographic activity, 4 EMG2-R wireless modules from Biopac were used to collect the signal from the Bionomadix BN-TX model Biopac transmission modules
- The RX130B electrogoniometer from Biopac was used to record the goniometry
- For the registration of the reaction force of the soil, the piezoelectric platform of forces AMTI brand, USA model SGA6-4 was used
- The surface electrodes used were bipolar (Ag/Ag Cl Blue Sensor N-00-5 Medicotest)
- Abrasive gel, conductive gel, alcohol and cotton were used to prepare the skin
- Hypoallergenic adhesive (Hypafix 10 cm × 10 m) was used for electrogoniometer fixation
- The validity and reliability of the instruments used is assumed and proven by the manufacturer and his technical department. On the other hand, these instruments have been used in different studies to obtain the results required.⁷⁰⁻⁷⁴

Study design

The type of study carried out is an observational study in which, using surface electromyography (EMG), a force platform and an electrogoniometer, the aim is to assess muscle activation and its relationship with GRF (specifically the vertical component F_z), having as main objective to analyze the correlation of activation of the different muscles and the ground reaction force (GRF) detected.

The subjects who have participated in the study are healthy subjects, both sexes, aged between 18 and 30 years and without any pathology or disease that can influence on the balance. Subjects that report an ACL lesion have also been ruled out.

They have also been excluded from the study all those subjects who report having some type of injury or sequel that does not allow them to perform a jump normally or that the jump may be contraindicated with their injury or pathology. Therefore, the study included 74 subjects of which 35 (47.3%) were women and 39 (52.7%) men, with a median age of 23.1 years, a median height of 1.73 m and a median weight of 69.37 kg (see descriptive data of the study population in [Table 1](#)).

Selection criteria

The inclusion criteria for incorporating subjects into the study were an age between 18 and 30 years old, both genders, being healthy and the acceptance in participating in the study by signing the informed consent. The exclusion criteria were being pregnant, having any pathology or disease that can influence on balance, previous or current ACL injury, having some type of injury or sequel that does not allow them to perform a normal jump or any contraindication of jumping.

Ethical considerations

The approval for the completion of this study was requested from the Ethics and Research Commission of the Blanquerna School of Psychology, Education and Sports Sciences of the Ramon Llull University. The favorable resolution was obtained on January 17, 2018.

To the subjects, once their participation was accepted and the informed consent signed, a small file was opened in which the personal data were recorded, as well as relevant data for the study (history and clinical history). Once these data were collected, the skin was prepared where the EMG electrodes were to be placed (cleaning and shaving if necessary). The electrodes were as described by Cram and Casman⁶⁹ in the following muscle groups: quadriceps (vastus medialis and vastus lateralis); hamstrings (semimembranosus/semitendinosus and biceps femoris); tibialis anterior; gastrocnemius; gluteus maximus; gluteus medius.

Likewise, the electronic goniometer was placed on the dominant limb, following the alignment of the limb and adhering it to the skin by a hypo-allergenic adhesive plaster. Once the electrodes and electrogoniometer were placed, a check was made to ensure that the receivers emitted information and from that moment a new calibration of the electrogoniometer was performed, in supine position and sitting; also, the goniometer was calibrated prior to use. Once the initial checks had been made, the baseline data (supine and standing) and maximum contraction values of the groups involved were acquired. The acquisition of the maximum contraction values was made and the subjects were placed on the force platform and their static position was assessed in the 3 axes.

For the election of the dominant side, subjects were asked to kick a ball as they normally would do (they were not given more information) and introduce it between two cones separated from each other by 1.5 m and located 4 m. The leg chosen for the kick would be considered dominant.

Once this data was collected, we start with the jumps. For this, the subjects were asked to climb on a 45 cm high stool and located 35 cm away from the edge of the platform. From this initial position and following the indications of the research team, the subjects were asked to jump with both legs on the platform and as soon as they touched the platform, make a vertical jump (the largest possible that would allow them to stabilize later) and land on his dominant leg and stabilize the position, keeping it until they were told it was necessary (see [Fig. 1](#) and [Table 2](#)).

For this, the necessary number of jumps were made until 3 good jumps could be obtained. They were considered

Table 1 Descriptive data of the study population.

	Men	Women
Gender <i>n</i> (%)	35 (47.3%)	39 (52.7%)
Age (median)	24.2 years	21.69 years
Height (median)	1.73 m	1.61 m
Weight (median)	79.38 kg	58.78 kg
Physically active	82.05%	77.14%




Figure 1 Description of the jump and landing.

Table 2 Description of the moments of the jump and landing.

Acronym	Description
P0	Flight start instant ($F_v = 0$)
P4	Ending instant of the flight ($F_v < 0$)
P1	Flight midpoint = $((P_4 - P_0)/2) + P_0$
P2	1/3 time between P1 and P4 = $P_1 + 1/3*(P_4 - P_1)$
P3	2/3 time between P1 and P4 = $P_1 + 2/3*(P_4 - P_1)$
P5	P4 + 30 ms
P6	P4 + 40 ms
P7	P4 + 70 ms
P8	P4 + 100 ms
P9	Instant in which F_v is maximum in the reception of the jump in the equilibrium phase
P10	Midpoint between P9 and P11 = $P_9 + (P_{11} - P_9)/2$
P11	Instant when the knee flexion-extension angle of the dominant leg (Gonio X) is maximum

good, all those that the subjects were able to stabilize, landed inside the platform and were unipodal.

Statistical analysis

A descriptive analysis was carried out for all the variables collected in the DCN (data collection notebook). Prior to the analysis, the Shapiro Wilks and Kolmogorov-Smirnov tests were performed together with the QQplot graphs to determine the normality of the variables studied. For the majority of the data, clear violations of the normality in the distribution of the data were observed, so it was decided to work according to a non-normal distribution. To study the relationship between variables, the non-parametric linear correlation between variables was used using the Spearman correlation coefficient statistic. The analyzes were performed based on the available dates, without using absent value substitution techniques, and describing the number of missing data for each analysis. In all statistical tests performed a significance level of 5% was used (P

value = 0.05). All analyzes were carried out with the SPSS v.25 program.

Results

Once the results obtained have been analyzed, significant differences have been observed in P9–P11 interval. The P -value, it always been <0.001 . As can be seen in Table 3, at instant P11, there is a strong correlation (R) between the reaction force of the soil (F_z) with average activation between instants P9–11 of the vastus lateralis ($R=0.511$) and tibialis anterior ($r=0.539$) muscles. In the case of women, at instant P11, there is a strong correlation (R) between the reaction force of the soil (F_z) with average activation between instants P9–11 of the biceps femoris ($R=0.545$), vastus lateralis ($R=0.583$), tibialis anterior ($R=0.522$) and gluteus maximus ($R=0.529$). On the other hand, in relation to the ground reaction force at instant P11, there is a strong correlation (R) with the activation averages of the vastus medialis ($R=0.670$), tibialis anterior ($R=0.569$), gluteus maximus ($R=0.649$) and gluteus medius ($R=0.571$) muscles.

In the rest of the moments or intervals of the jump and landing, no significant differences were found.

Discussion

The muscular activity and the correlation in the activation of the different muscles, is an aspect that has been commented by many authors since it is considered important for the prevention of ACL injuries. These muscle correlations tell us about the ability to “work together” and to achieve a good body stabilization after an activity like as in this case, the landing after a jump and to counteract the ground reaction forces (GRF) and thus avoid or decrease a potential risk of injury. There are different authors who point out that there are biomechanical profiles that relate high GRF values with little damping and therefore with a higher risk of injury.

In this sense, the results obtained show that, during the entire jump and landing phase, there are have only found significant differences in the period P9–P11 being P9 the point where the reaction force of the ground is greater and P11 the moment where the knee reaches the maximum flexion after landing. The fact that the differences have occurred in those instants and intervals, supports the

Table 3 Correlation between Fz and muscle activation in the P9P11 interval.

	FzP11								
	Women			Men			Women & men		
	R	P-value	N	R	P-value	N	R	P-value	N
BF_Pr_P9P11	,545	0.001	35						
VL_Pr_P9P11	,583	<0.001	35				,511	<0.001	74
TA_Pr_P9P11	,522	0.001	35	,569	<0.001	39	,539	<0.001	74
GMa_Pr_P9P11	,529	0.001	35	,649	<0.001	39			
VM_Pr_P9P11				,670	<0.001	39			
GMe_Pr_P9P11				,571	<0.001	39			

Acronyms and abbreviations:

BF: electromyographic value of the biceps femoris.

Fz: vertical component of the ground reaction force.

GMa: electromyographic value of the gluteus maximus.

GMe: electromyographic value of the gluteus medius.

P9–P11: interval between P9 and P11.

R: correlation coefficient.

TA: electromyographic value of the tibialis anterior.

VL: electromyographic value of the vastus lateralis.

VM: electromyographic value of the vastus medialis.

idea that, beyond the instants in which the different authors point out that in which the injury normally occurs (interval between P6 and P8), the neuromuscular activity and co-contractions continue to stabilize the joint and allow the individual to continue with the activity or the consequent movements that will occur.^{69–77} The significance of the practical application of the results is that these results will allow to evaluate in more detail these phases of the landing and give more importance to the late stabilization phase after a jump, giving value to this final phase of stabilization, which even if it is not the phase in which the injury usually occurs, it is necessary to chain activities that occur in sports practice; normally the activities are continued activities, not isolated actions. And so, it is important a good stabilization after each action and that will allow to continue with the activity without injury. And this phase of the landing will be important to consider when planning prevention work as well as when the patient is being rehabilitated.

Regarding with this, it can be noted that there is a high correlation of the activation of certain muscle groups with the GRF in P11; in the case of women, there is a strong correlation with that point with the activity of the biceps femoris, vastus lateralis, tibialis anterior and gluteus maximus muscles at that point P11, while in the case of men, the muscles are vastus lateralis, tibialis anterior and gluteus medius. The muscles of the posterior chain have been previously indicated as essential for the stabilization and absorption of GRF.⁵⁴ And of course, the co-contraction with the muscles of the anterior chain is necessary for the joint stabilization. On the other hand, also shows the differences that previously have been mentioned in results, in which it is been highlighted the greater activation by women in the gluteus maximus, and the gluteus medius being more active in men. This also agrees with what other authors have indicated; that the mechanisms and reasons for injury may be different depending on sex.³¹ This will be important to consider when planning the training and prevention exercises, because this could help to reduce the incidence of injuries.

These results lead to conclude that, as other authors point out, depending on sex, there are different neuromuscular mechanisms to face an action such as landing after a jump and this justifies the importance of knowing what are the patterns of activation to prevent an important injury as the ACL injury is but not only in the moment where the injury occurs. It is also important to point out the different use that the subjects do of the gluteus maximus and gluteus medius (two very important muscles for their stabilizing and locomotion capacity), depending on the sex, in this late stage of stabilization.

As a possible methodological limitation of the study, it should be noted that the possible muscular or psychological fatigue that the subject presented by activities done in the previous days, was not considered.

Therefore, for future studies, the fatigue will be considered, because it is believed that it can play an important role in neuromuscular patterns. And of course, it is considered important to study and analyze in more detail the differences in neuromuscular pattern that may occur due to sex.

In conclusion, the evaluation of GRF and muscle activation patterns, allows to assess the dynamics of landing after a jump and to be able to detect different patterns according to sex, with the consequent importance that it can have in the injury mechanism.

Conflict of interest

The authors declare that they don't have any conflict of interest.

Uncited references

Refs. 46–51, 53, 54.

References

- Beutler A, de la Motte S, Marshall S, Padua D, Boden B. Muscle strength and qualitative jump-landing differences in male and female military cadets: the jump-acl study. *J Sports Sci Med*. 2009;8:663–71.
- Webster KA, Gribble PA. Time to stabilization of anterior cruciate ligament-reconstructed versus healthy knees in National Collegiate Athletic Association Division I female athletes. *J Athl Train*. 2010;45:580–5.
- Lindenberg KM, Carcia CR, Phelps AL, Martin RL, Burrows AM. The influence of heel height on sagittal plane knee kinematics during landing tasks in recreationally active and athletic collegiate females. *Int J Sports Phys Ther*. 2011;6:186–98.
- Shultz SJ, Nguyen AD, Leonard MD, Schmitz RJ. Thigh strength and activation as predictors of knee biomechanics during a drop jump task. *Med Sci Sports Exerc*. 2009;41:857–66.
- Hanson AM, Padua DA, Troy Blackburn J, Prentice WE, Hirth CJ. Muscle activation during side-step cutting maneuvers in male and female soccer athletes. *J Athl Train*. 2008;43:133–43.
- Schmitz RJ, Shultz SJ. Contribution of knee flexor and extensor strength on sex-specific energy absorption and torsional joint stiffness during drop jumping. *J Athl Train*. 2010;45:445–52.
- Shultz SJ, Schmitz RJ. Effects of transverse and frontal plane knee laxity on hip and knee neuromechanics during drop landings. *Am J Sports Med*. 2009;37:1821–30.
- Shultz SJ, Schmitz RJ, Nguyen AD, Levine BJ. Joint laxity is related to lower extremity energetics during a drop jump landing. *Med Sci Sports Exerc*. 2010;42:771–80.
- Lephart SM, Abt JP, Ferris CM, Sell TC, Nagai T, Myers JB, et al. Neuromuscular and biomechanical characteristic changes in high school athletes: a plyometric versus basic resistance program. *Br J Sports Med*. 2005;39:932–8.
- Louw Q, Grimmer K, Vaughan C. Knee movement patterns of injured and uninjured adolescent basketball players when landing from a jump: a case-control study. *BMC Musculoskel Disord*. 2006;7:22.
- Schmitz RJ, Shultz SJ, Nguyen AD. Dynamic valgus alignment and functional strength in males and females during maturation. *J Athl Train*. 2009;44:26–32.
- Hägglund M, Atroshi I, Wagner P, Waldén M. Superior compliance with a neuromuscular training programme is associated with fewer ACL injuries and fewer acute knee injuries in female adolescent football players: secondary analysis of an RCT. *Br J Sports Med*. 2013;47:974–9.
- McLean SG, Walker K, Ford KR, Myer GD, Hewett TE, van den Bogert AJ. Evaluation of a two dimensional analysis method as a screening and evaluation tool for anterior cruciate ligament injury. *Br J Sports Med*. 2005;39:355–62.
- Myer GD, Ford KR, Brent JL, Hewett TE. Differential neuromuscular training effects on ACL injury risk factors in “high-risk” versus “low-risk” athletes. *BMC Musculoskel Disord*. 2007;8:39.
- Sasaki S, Nagano Y, Ichikawa H. Loading differences in single-leg landing in the forehand- and backhand-side courts after an overhead stroke in badminton: a novel tri-axial accelerometer research. *J Sports Sci*. 2018;1–8.
- Leppänen M, Pasanen K, Kujala UM, Vasankari T, Kannus P, Äyrämö S, et al. Stiff landings are associated with increased ACL injury risk in young female basketball and floorball players. *Am J Sports Med*. 2017;45:386–93.
- Weiss K, Whatman C. Biomechanics associated with patellofemoral pain and ACL injuries in sports. *Sport Med*. 2015;45:1325–37.
- Donnell-Fink LA, Klara K, Collins JE, Yang HY, Goczalk MG, Katz JN, et al. Effectiveness of knee injury and anterior cruciate ligament tear prevention programs: a meta-analysis. *PLoS One*. 2015;10, e0144063.
- Gokeler A, Neuhaus D, Benjaminse A, Grooms DR, Baumeister J. Principles of motor learning to support neuroplasticity after ACL injury: implications for optimizing performance and reducing risk of second ACL injury. *Sport Med*. 2019;49:853–65.
- Dai B, Garrett WE, Gross MT, Padua DA, Queen RM, Yu B. The effects of 2 landing techniques on knee kinematics, kinetics, and performance during stop-jump and side-cutting tasks. *Am J Sports Med*. 2015;43:466–74.
- Gokeler A, Benjaminse A, Welling W, Alferink M, Eppinga P, Otten B. The effects of attentional focus on jump performance and knee joint kinematics in patients after ACL reconstruction. *Phys Ther Sport*. 2015;16:114–20.
- Kotsifaki A, Korakakis V, Whiteley R, Van Rossom S, Jonkers I. Measuring only hop distance during single leg hop testing is insufficient to detect deficits in knee function after ACL reconstruction: a systematic review and meta-analysis. *Br J Sports Med*. 2019, bjsports-2018-099918.
- Nessler T, Denney L, Sampley J. ACL injury prevention: what does research tell us? *Curr Rev Musculoskel Med*. 2017;10:281–8.
- Sugimoto D, Myer GD, Barber Foss KD, Pepin MJ, Micheli LJ, Hewett TE. Critical components of neuromuscular training to reduce ACL injury risk in female athletes: meta-regression analysis. *Br J Sports Med*. 2016;50:1259–66.
- Silvers-Granelli HJ, Bizzini M, Arundale A, Mandelbaum BR, Snyder-Mackler L. Does the FIFA 11 + injury prevention program reduce the incidence of acl injury in male soccer players? *Clin Orthop Relat Res*. 2017;475:2447–55.
- Filbay SR, Grindem H. Evidence-based recommendations for the management of anterior cruciate ligament (ACL) rupture. *Best Pract Res Clin Rheumatol*. 2019.
- Horvath A, Senorski EH, Westin O, Karlsson J, Samuelsson K, Svantesson E. Outcome after anterior cruciate ligament revision. *Curr Rev Musculoskel Med*. 2019;12:397–405.
- Nyman E, Armstrong CW. Real-time feedback during drop landing training improves subsequent frontal and sagittal plane knee kinematics. *Clin Biomech*. 2015;30:988–94.
- Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *Br J Sports Med*. 2016;50:804–8.
- Larsen JB, Farup J, Lind M, Dalgas U. Muscle strength and functional performance is markedly impaired at the recommended time point for sport return after anterior cruciate ligament reconstruction in recreational athletes. *Hum Mov Sci*. 2015;39:73–87.
- Pfeifer CE, Beattie PF, Sacko RS, Hand A. Risk factors associated with non-contact anterior cruciate ligament injury: a systematic review. *Int J Sports Phys Ther*. 2018;13:575–87.
- DeMorat G, Weinhold P, Blackburn T, Chudik S, Garrett W. Aggressive quadriceps loading can induce noncontact anterior cruciate ligament injury. *Am J Sports Med*. 2004;32:477–83.
- Boden BP, Dean GS, Feagin JA, Garrett WE. Mechanisms of anterior cruciate ligament injury. *Orthopedics*. 2000;23:573–8.
- Boden BP, Griffin LY, Garrett WE. Etiology and prevention of noncontact ACL injury. *Phys Sportsmed*. 2000;28:53–60.
- Olsen OE, Myklebust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med*. 2004;32:1002–12.
- Pflum MA, Shelburne KB, Torry MR, Decker MJ, Pandy MG. Model prediction of anterior cruciate ligament force during drop-landings. *Med Sci Sports Exerc* [Internet]. 2004;36:1949–58.
- Urabe Y, Kobayashi R, Sumida S, Tanaka K, Yoshida N, Nishiwaki GA, et al. Electromyographic analysis of the knee during jump landing in male and female athletes. *Knee*. 2005;12:129–34.

38. Shultz SJ, Schmitz RJ, Benjaminse A, Collins M, Ford K, Kulas AS. ACL research retreat VII: an update on anterior cruciate ligament injury risk factor identification screening, and prevention. *J Athl Train.* 2015;50:1076–93.
39. Bakker R, Tomescu S, Breneman E, Hangalur G, Laing A, Chandrashekar N. Effect of sagittal plane mechanics on ACL strain during jump landing. *J Orthop Res.* 2016;34:1636–44.
40. Blackburn JT, Padua DA. Sagittal-plane trunk position, landing forces, and quadriceps electromyographic activity. *J Athl Train.* 2009;44:174–9.
41. Shimokochi Y, Yong Lee S, Shultz SJ, Schmitz RJ. The relationships among sagittal-plane lower extremity moments: implications for landing strategy in anterior cruciate ligament injury prevention. *J Athl Train.* 2009;44:33–8.
42. Kulas A, Zalewski P, Hortobagyi T, DeVita P. Effects of added trunk load and corresponding trunk position adaptations on lower extremity biomechanics during drop-landings. *J Biomech.* 2008;41:180–5.
43. Husted RS, Bencke J, Hölmich P, Andersen LL, Thorborg K, Bandholm T, et al. Maximal hip and knee muscle strength are not related to neuromuscular pre-activity during sidcutting maneuver: a cross-sectional study. *Int J Sports Phys Ther.* 2018;13:66–76.
44. Dyhre-Poulsen P, Krogsgaard MR. Muscular reflexes elicited by electrical stimulation of the anterior cruciate ligament in humans. *J Appl Physiol.* 2000;89:2191–5.
45. Zebis MK, Bencke J, Andersen LL, Døssing S, Alkjaer T, Magnusson SP, et al. The effects of neuromuscular training on knee joint motor control during sidcutting in female elite soccer and handball players. *Clin J Sport Med.* 2008;18:329–37.
46. Markolf KL, Gorek JF, Kabo JM, Shapiro MS. Direct measurement of resultant forces in the anterior cruciate ligament. An in vitro study performed with a new experimental technique. *J Bone Joint Surg Am.* 1990;72:557–67.
47. Meyer EG, Haut RC. Anterior cruciate ligament injury induced by internal tibial torsion or tibiofemoral compression. *J Biomech.* 2008;41:3377–83.
48. Hewett TE, Stroupe AL, Nance TA, Noyes FR. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24:765–73.
49. Krosshaug T, Nakamae A, Boden BP, Engebretsen L, Smith G, Slauterbeck JR, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2007;35:359–67.
50. Hewett TE, Myer GD, Ford KR, Paterno MV, Quatman CE. Mechanisms, prediction, and prevention of ACL injuries: cut risk with three sharpened and validated tools. *J Orthop Res.* 2016;34:1843–55.
51. Kiapour AM, Demetropoulos CK, Kiapour A, Quatman CE, Wordeman SC, Goel VK, et al. Strain response of the anterior cruciate ligament to uniplanar and multiplanar loads during simulated landings. *Am J Sports Med.* 2016;44:2087–96.
52. Hewett TE, Zazulak BT, Myer GD, Ford KR. A review of electromyographic activation levels, timing differences, and increased anterior cruciate ligament injury incidence in female athletes. *Br J Sports Med.* 2005;39:347–50.
53. Hewett TE, Myer GD, Ford KR, Heidt RS, Colosimo AJ, McLean SG, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33:492–501.
54. Färber S, Heinrich D, Werner I, Federolf P. Is it possible to voluntarily increase hamstring muscle activation during landing from a snow jump in alpine skiing? – a pilot study. *J Sports Sci.* 2018;18:1–8.
55. Horita T, Komi PV, Nicol C, Kyröläinen H. Interaction between pre-landing activities and stiffness regulation of the knee joint musculoskeletal system in the drop jump: implications to performance. *Eur J Appl Physiol.* 2002;88:76–84.
56. Duncan A, McDonagh MJ. Stretch reflex distinguished from pre-programmed muscle activations following landing impacts in man. *J Physiol.* 2000;526 Pt 2:457–68.
57. McDonagh MJN, Duncan A. Interaction of pre-programmed control and natural stretch reflexes in human landing movements. *J Physiol [Internet].* 2002;544 Pt 3:985–94.
58. Cowling EJ, Steele JR, McNair PJ. Effect of verbal instructions on muscle activity and risk of injury to the anterior cruciate ligament during landing. *Br J Sports Med.* 2003;37:126–30.
59. McNair PJ, Prapavessis H, Callender K. Decreasing landing forces: effect of instruction. *Br J Sports Med.* 2000;34:293–6.
60. Prapavessis H, McNair PJ. Effects of instruction in jumping technique and experience jumping on ground reaction forces. *J Orthop Sports Phys Ther.* 1999;29:352–6.
61. Labanca L, Laudani L, Mariani P, Macaluso A. Postural adjustments following ACL rupture and reconstruction: a longitudinal study. *Int J Sports Med.* 2018;39:549–54.
62. Bennis N, Roby-Brami A, Dufossé M, Bussel B. Anticipatory responses to a self-applied load in normal subjects and hemiparetic patients. *J Physiol Paris.* 1996;90:27–42.
63. Kanekar N, Aruin AS. The effect of aging on anticipatory postural control. *Exp Brain Res.* 2014;232:1127–36.
64. Labanca L, Laudani L, Casabona A, Menotti F, Mariani PP, Macaluso A. Early compensatory and anticipatory postural adjustments following anterior cruciate ligament reconstruction. *Eur J Appl Physiol.* 2015;115:1441–51.
65. Bulgheroni P, Bulgheroni MV, Andriani L, Guffanti P, Giughello A. Gait patterns after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 1997;5:14–21.
66. Ferber R, Osternig LR, Woollacott MH, Wasielewski NJ, Lee JH. Gait perturbation response in chronic anterior cruciate ligament deficiency and repair. *Clin Biomech (Bristol, Avon).* 2003;18:132–41.
67. Lustosa LP, Ocarino JM, de Andrade MAP, Pertence AE, Bitencourt NF, Fonseca ST. Muscle co-contraction after anterior cruciate ligament reconstruction: Influence of functional level. *J Electromyogr Kinesiol.* 2011;21:1050–5.
68. Paterno MV, Schmitt LC, Ford KR, Rauh MJ, Myer GD, Huang B, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med.* 2010;38:1968–78.
69. Criswell E. *Cram's introduction to surface electromyography.* Jones & Bartlett Learning; 2010.
70. Oliver G, Portabella F, Hernandez JA. A comparative study of the neuromuscular response during a dynamic activity after anterior cruciate ligament reconstruction. *Eur J Orthop Surg Traumatol.* 2019;29:633–8.
71. Flaxman TE, Alkjaer T, Smale KB, Simonsen EB, Krogsgaard MR, Benoit DL. Differences in EMG-moment relationships between ACL-injured and uninjured adults during a weight-bearing multidirectional force control task. *J Orthop Res.* 2019;37:113–23.
72. Zebis MK, Andersen LL, Brandt M, Myklebust G, Bencke J, Lauridsen HB, et al. Effects of evidence-based prevention training on neuromuscular and biomechanical risk factors for ACL injury in adolescent female athletes: a randomised controlled trial. *Br J Sports Med.* 2016;50:552–7.
73. Suarez T, Laudani L, Giombini A, Saraceni VM, Mariani PP, Pigozzi F, et al. Comparison in joint-position sense and muscle coactivation between anterior cruciate ligament-deficient and healthy individuals. *J Sport Rehab.* 2016;25:64–9.
74. Davis K, Williams JL, Sanford BA, Zucker-Levin A. Assessing lower extremity coordination and coordination variability in individuals with anterior cruciate ligament reconstruction during walking. *Gait Posture.* 2019;67:154–9.

75. Heebner NR, Rafferty DM, Wohleber MF, Simonson AJ, Lovalekar M, Reinert A, et al. Landing kinematics and kinetics at the knee during different landing tasks. *J Athl Train.* 2017;52:1101–8.
76. Christoforidou A, Patikas DA, Bassa E, Paraschos I, Lazaridis S, Christoforidis C, et al. Landing from different heights: biomechanical and neuromuscular strategies in trained gymnasts and untrained prepubescent girls. *J Electromyogr Kinesiol.* 2017;32:1–8.
77. Huurnink A, Fransz DP, Kingma I, de Boode VA, Dieën JHvan. The assessment of single-leg drop jump landing performance by means of ground reaction forces: a methodological study. *Gait Posture.* 2019;73:80–5.